

CATALOGUE VETERINARY DIVISION

www.ginyechina.com



山东金野生物科技有限公司
Shandong Ginye Biotech Co., Ltd

FOR ANIMAL · FOR HEALTH · FOR YOU



CONTENT

About Ginye	03
Nutritional Feed Supplement	04-18
Herbal Extract	19-32
Antibiotic	33-62
Products List	63-64



GINYE DEVELOPMENT HISTORY



Shandong Ginye Biotech Co., Ltd. is a integrated company whose main business is animal pharmaceutical / feed additive / aquatic products / pet products research development, production and trading.

With 2 modern GMP factories (10 production lines) to produce more than 200 products. Our products include: powder / oral solution / liquid injection / tablet / injectable powder / granula / herbal extract / disinfectant / pesticide.





GINVITA W/S

Multivitamin + Minerals+ Amino Acids WSP

Nutrition

For Veterinary Use Only

Composition: Each kg contains

Vit A	5,000,000 IU	Vit D3.....	2,000,000 IU
Vit E.....	15,000 mg	Vit B1	3,000 mg
Vit B2.....	5,000 mg	Vit B6.....	8,000 mg
VitB12.....	50 mg	Vit C	50,000 mg
Vit K3	500 mg	Folic Acid	1,000 mg
Biotin	50 mg	Choline chloride.....	10,000 mg
Manganese sulphate:	2000mg	Magnesium sulphate:	2000mg
Zinc sulphate:	2000mg	Iron sulphate:	1500mg
Lysine:	15,000mg	Methionine:	12,000mg

Description:

- Quickly stress reliever
- Stimulates antibodies production for better immunity.
- Vaccine will be utilized properly and completely.
- Improve egg production and maintains good persistency.
- Broilers become more active with good growth and body weights.

Usage & Dosage:

For oral administration

Poultry swine: 1g per 2-3 liter water continue for 3-5 days

Calves, goats and sheep : 1 g per 10 kg. body weight for 3 - 5 days.

Cattle : 1 gram per 20 kg. body weight for 3 - 5 days.

Doubled when use mixed with feed.

Medicated drinking water should be used within 24 hours.

Withdrawal Period:

None

Storage :

Store in cool place prevent from light.

Keep out of reach of children.

Packing :

1kg/Bag in Bucket or Carton

Validity :

3 years



GINYE BIOTECH EXPORT

GIN-TOX W/S

Mycotoxin Cleaner, Detoxification, Immune
Booster Poultry Formula

Composition per gram:

Each gram contains

Yeast Polysaccharide. $\geq 50\text{mg}$

Bacillus subtilis $\geq 1 \times 10^9 \text{ CFU}$

Carrier : Glucose oxidase Potassium sorbate ,
Herb extract etc.

Description:

1. Degrade mycotoxin and glucose organically, consume oxygen in intestinal tract, increase acid environment. Restrain harmful bacteria reproduction, decrease intestinal tract disease.
2. Reduce feed mildew obviously, restrain all kinds of mycotoxin, promote body toxin metabolism.
3. Enhance digestive enzyme activity, improve appetite, promote feed digestion, increase feed conversion.
4. Improve immunity and liver kidney detoxification function, it has good detoxification for disease and medicine poisoning.

Usage & Dosage:

1 g per 3 liter of drinking water keep more than 7 days.
Double dosage during mycotoxin serious period.

Withdrawal Period:

None

Storage :

Stored with sealed packaging in a cool, dry and ventilated place.
Avoid to be exposed to the sun, rain, high temperature and high humidity.

Packing :

500g 1kg bags other bulk packing forms.

Validity :

3 years

Nutrition

For Veterinary Use Only





TOXIMON PLUS

Upgrade Formula Mycotoxin Binder

Nutrition

For Veterinary Use Only

Composition:

Modified hydrate sodium calcium aluminosilicate	40%
Organic polysaccharide montmorillonite	30%
Yeast cell wall	30%

Function:

- 1.Strongly adsorb mycotoxin in feed and animal body, not affect the absorption of nutrients in the body.
- 2.Remove immune suppression and improve animal immunity.
- 3.Protect intestinal mucosa, repair and improve the defense function of the intestinal mucosal barrier.
- 4.Prevent and control diarrhea ensure normal digestion and absorption of various nutrients.

Usage & Dosage:

Prevent mycotoxins: 0.5-1.5kg/ton
Adsorb and remove mycotoxins: 1.5-4kg/ton
Livestock : 1-2kg/ ton
Aquaculture: 2-4kg/ton
Additive amount according to mycotoxin contamination situation, or consult with veterinarian

Withdrawal Period:

None

Storage :

Should be stored in ventilated, dry warehouse, stacking should be away from the ground and wall, strictly prohibit mixing with toxic, harmful substances.

Packing :

25kg Polypropylene (PP) bag

Validity :

3 years



GINYE BIOTECH EXPORT

GINVITA PLUS

Multi-Function Nutrition Supplement Poultry Formula

Nutrition

For Veterinary Use Only



Composition: each gram contains

Astragalus Extract.....	100 mg	Taurine.....	100 mg
β glucan.....	25 mg	Vitamin A.....	10000 IU
Vitamin D3.....	5000 IU	Vitamin E.....	20 mg
Vitamin B1	5 mg	Vitamin B2.....	8 mg
Vitamin B6.....	8 mg	Vitamin B12.....	50 mcg
Vitamin C.....	30 mg	Vitamin K3.....	5 mg
Nicotinamide.....	60 mg	Folic acid.....	2 mg
Biotin.....	300 µg	Choline chloride.....	10 mg
L-Lysine.....	50 mg	DL-Methionine.....	45 mg

Description:

1. Adjust intestinal flora balance, decrease enteritis and diarrhea diseases.
2. Multivitamin supplementation, keep poultry physiological function,
3. improve immunity and anti-stress strength, decrease mortality.
4. Stomachic attractant and promote digestion & ingestion, improve FCR.

Usage & Dosage:

Poultry: 1 kg per 1500-2000 Liter water keep 3-5 days
Double dosage during disease and recovery period.
Caution: This product cannot mix use with other medicine and vaccine, use interval time should not less than 3 hours.

Storage :

Store in cool place prevent from light.
Keep out of reach of children.

Packing :

250g 1kg 5kg Bags

Validity :

3 years





GINEGG PREMIX

Natural Herb Extract Probiotics Egg Promoter

Nutrition

For Veterinary Use Only

Composition: Each gram contains

Astragalus polysaccharide & Zymosan.....100 mg

Bacillus subtilis $\geq 5 \times 10^9$ CFU

Carrier: upto 1 g

Indications:

- Improve eggshell hardness decrease broken egg rate .
- Help Layer establish good immune system, little disease happened.
- Extend laying fastigium Increase good quality egg production .
- Strengthen liver and kidney function, strengthen anti-stress capability.
- Keep intestinal tract microecosystem balance, decrease digestive system disease, Reduce feed & egg ratio.

Dosage&Administration:

1. Use during laying early period (about 60days) and laying late stage (physical laying rate drop off period). 1kg per 1000kg feed.
2. Double dosage during stress & disease recovery period, use more than 10 days. Medicated feed should be used within 3-4 hours, better use it during sunset.

Withdrawal Period:

None

Storage :

Keep in storage below 30°C, prevent from light.

Packing :

1kg×15bags/carton 25kg

Validity :

3 years



GINYE BIOTECH EXPORT

GINEGG ORAL

Natural Herb Extract Egg Promoter

Nutrition

For Veterinary Use Only



Composition: Each ml contains

Astragalus polysaccharide & Zymosan.....100 mg
Soybean oil, Glucose oxidase other solvents upto..... 1 ml

Indications:

- Improve eggshell hardness decrease broken egg rate .
- Help Layer establish good immune system, little disease happened.
- Extend laying fastigium Increase good quality egg production .
- Strengthen liver and kidney function, strengthen anti-stress capability.
- Keep intestinal tract microecosystem balance, decrease digestive system disease, Reduce feed & egg ratio.

Dosage&Administration:

1. Use during laying early period (about 60days) and laying late stage (physical laying rate drop off period).1kg per 1000kg feed.
2. Double dosage during stress & disease recovery period, use more than 10 days.
Medicated feed should be used within 3-4 hours, better use it during sunset.

Withdrawal Period:

None

Storage :

Store in a dry, dark place between 5°C and 25°C.
Keep it away from children.

Packing :

1kg×15bags/carton 25kg

Validity :

3 years





GINVITA ORAL

Multivitamin + Amino Acids Solution

Nutrition

For Veterinary Use Only

Composition: each gram contains

Vit A	8,000 IU	Vit D3.....	4,000 IU
Vit E.....	20 mg	Vit B1	3 mg
Vit B2.....	5 mg	Vit B6.....	5 mg
VitB12.....	50µg	Vit C.....	30 mg
Vit K3.....	10mg	Folic Acid	1 mg
Biotin	50µg	D-panthenol	20 mg
Nicotinamide.....	25 mg	DL-Methionine	6 mg
Lysine	10 mg	Glycine	3 mg
L-Arginine.....	2 mg	L-Threonine.....	1 mg
25-hydroxyvitamin D..	30mg	Xylooligosaccharide..	8 mg

Description:

GINVITA ORAL is a well balanced combination of essential vitamins and amino acid 25-hydroxyvitamin D for farm animals.

Prevention or treatment of vitamin deficiencies in farm animals.

Prevention or treatment of stress (caused by vaccination, diseases, transport, high humidity, high temperatures or extreme temperature changes).

Improvement of feed conversion.

Usage & Dosage:

Poultry: For broiler all stage and Layer brooding period

1ml per 2.5 L - 3 L drinking water keep 5-7 days or through the breeding period.

Double dosage during anti-stress and disease recovery period.

Livestock:

Calves, goats and sheep: 1 ml per 10 kg body weight for 3 - 5 days.

Cattle: 1 ml per 20 kg body weight for 3 - 5 days.

Swine: 1 litre per 1000 litre drinking water for 3 - 5 days.

Medicated drinking water should be used within 12 hours.

Storage :

Store in a dry, dark place between 5°C and 25°C. Keep it away from children.

Packing :

1L×12Bottle/CTN 500ML×30Bottle/CTN

Validity :

3 years



GINYE BIOTECH EXPORT

GIN-ACID ORAL

Complex Organic Acids Acidifier

Nutrition

For Veterinary Use Only



Composition: each liter contains

Formic acid 150g, Acetic acid 150g Lactic acid 100g , Propionic acid 20g, Malic Acid 10g, Other short chain fatty acids and their salts 100g.

Indications:

1. Effective against pathogenic bacteria like E. coli, Salmonella etc.
2. Improving protein digestion and FCR.
3. Stimulates the production of digestive enzymes
4. Prevent drinking line and papilla blocking, bacteria & virus growth, clean organicsubstance and residual mineral in drinking system. suitable for your all kinds ofdrinking system.
5. Reduces diarrhea and mortality rates

Dosage&Administration:

Drinking water hygiene : 0.1%-0.2% regular intervals drinking water, every 2-3 days.

Anti-Stress: 0.1%-0.2% drinking water, 2-3 days.

Nutritional diarrhea: 0.2%-0.4% drinking water, 3-5 days.

Water line cleaning : 1%-2% mix with water, steep more than 12 hours , wash it 2 times with clean water.

Withdrawal Period:

None

Storage :

Store in a dry, dark place between 5°Cand 25°C.
Keep it away from children.

Packing :

1L 5L 25L Plastic Container.

Validity :

3 years





GIN-GRO

Probiotic Feed Nutritional Additive
Ruminants Growth Promoter

Nutrition

For Veterinary Use Only

Composition: Each gram contains

Bacillus subtilis: 8×10^8 CFU/g

Lactobacillus: 2×10^8 CFU/g

Total Viable Bacteria: $\geq 1 \times 10^9$ CFU/g

Yeast Extract : 30 mg

Prebiotics (bifidus factor, oligosaccharide) Maifanite, Aspergillus oryzae culture

Protease Cellulase and other active enzyme upto 1g

Indications:

- 1.Improve rumen environment and buffer capacity, stabilize rumen pH value.
- 2.Promote rumen beneficial microorganism growth - Probiotics somatomedins provide nutrients to rumen beneficial microflora, its growth in number and vitality promptly.
- 3.Improve feed quality, promote feed digestion and absorption, adjust amino acid and fat metabolism improve meat quality.
- 4.Decrease digestion system disease, prevent diarrhea, Improve immunity, reduce ammonia in animal house.

Dosage&Administration:

Beef cattle : 3-5 kg/1000kg of feed

Sheep Goat : 2-3kg/1000kg of feed

Young animals: 2 kg/1000kg of feed

Dairy: 50g-100g/dairy cow/day

It is recommended to use this product through the whole feeding period. Dosage can be increased under disease condition.

Withdrawal Period:

None

Storage :

Keep in storage of 5-25°C,
prevent from air and light.
Keep it away from children.

Packing :

1kg 20kg Bag

Validity :

3 years



GINYE BIOTECH EXPORT

GIN-PRO ORAL

Probiotic Feed Nutritional Additive

Nutrition

For Veterinary Use Only



Composition: Each gram contains

Bacillus subtilis $\geq 2 \times 10^9$ cfu
Bacillus licheniformis $\geq 1 \times 10^9$ cfu
Enterococcus faecalis $\geq 2 \times 10^9$ cfu
Lactobacillus Bifidobacterium and other
beneficial bacteriums and their metabolite

Indications:

1. Improve disease resistance, Improve feed conversion rate, adjust gastrointestinal pH, improve intestinal environment and prevent diseases.
2. Reduce the occurrence of enteritis, prevent overfeeding and effectively prevent various types of coccidiosis in broiler.
3. Improve the egg production rate of layers, extend the peak period of egg production, increase the egg production rate in the middle and late stages,
4. Reduce the concentration of harmful gases such as ammonia and hydrogen sulfide in the animal house. improve the breeding environment and reduce pollution.

Dosage&Administration:

Poultry and rabbits : 1.5-2 ml/ 1 liter drinking water
cattle: 75-100 ml per animal.
calves: 30-50 ml per animal.
Sheep & goat: 20-30 ml per animal.

Withdrawal Period:

None

Storage :

Store in a dry, dark place between 5°C and 25°C.
Keep it away from children.

Packing :

500ml 1000ml bottle

Validity :

3 years





IMUGIN WS

Super Immunity Booster New Update Formula

Nutrition

For Veterinary Use Only

Composition: Each gram contains

Lysozyme	250 mg
Vitamin E	100 mg
Vitamin C	50 mg
Selenium	20 mg

Indications:

- Help animals immune organs growth promote get strong immunity.
- Stimulates antibodies production, release animals stress.
- Help vaccine be utilized properly and completely.
- Improve layer egg production and breeder fertility rate .
- Improve broilers production performance at lower cost.

Dosage&Administration:

For oral administration

1g per 2 Liter water continue for 5-7 days.

1g per 3 Liter of water for daily health care.

Other dosage should follow veterinarian advice.

Medicated drinking water should be used within 24 hours.

Withdrawal Period:

None

Storage :

Store in a dry, dark place below 30°C.

Keep it away from children.

Packing :

1kg/Bag 15kg/carton or bucket

Validity :

3 years



GINYE BIOTECH EXPORT

E-SE 100 ORAL

Oral Vitamin E For Veterinary Use

Nutrition

For Veterinary Use Only



Composition: Each ml contains

Vitamin E.....	100 mg
Sodium selenite.....	1.5 mg
Solvents ad	1 ml

Indications:

E-SE 100 Oral is indicated for vitamin E and/or selenium deficiency in calves, lambs, sheep, goats, piglets and poultry. Encephalo-malacia (Crazy Chick Disease), muscular dystrophy (White Muscle Disease, Stiff Lamb Disease), exudative diathesis (generalised oedematous condition), decreased hatchability of eggs.

Dosage&Administration:

For oral administration via drinking water.

Calves, lambs, sheep, goats, piglets : 10 ml per 50 kg bodyweight during 5 - 10 days.

Poultry : 1ml per 1.5-2 litre of drinking water during 5 - 10 days.

Medicated drinking water should be used within 24 hours.

Other dosage should comply with veterinarian's suggestion

Withdrawal Period:

None

Storage :

Store in dry dark place between 5°C and 25 °C,
Store in closed packing.

Packing :

In a 500ml 1L plastic bottle.

Validity :

3 years





LIVERGIN ORAL

Liver Tonic Oral Liquid

Nutrition

For Veterinary Use Only

Composition: Each ml contains

Sorbitol.....	200 mg
Carnitine HCL	25 mg
Choline Chloride.....	18.5 mg
DL-Methionine	10 mg
Magnesium	10 mg
Astragalus Extract.....	20 mg

Indications:

LIVERGIN is a metabolic enhancer & liver-protector which protects animal's liver detoxification, reducing clinical symptoms when feed is contaminated with pesticides, mycotoxins or after long antibiotic treatment. Prevention and treatment of Fatty Liver syndrome. Improve immunity and promoting animal growth.

Dosage&Administration:

For oral administration

Poultry: 1ml per 2-2.5L of drinking water

Doubled during serious condition.

Medicated drinking water should be used within 24 hours.

WITHDRAWAL PERIOD:

None

Storage :

Store in a dry, dark place between 5°C and 30°C.

Keep it away from children.

Packing :

1L 5L container

Validity :

3 years



GINYE BIOTECH EXPORT

GINFAT W/S

Growth Promoter Nutritional Additive

Nutrition

For Veterinary Use Only



Composition: Each gram contains

Taurine100 mg
Astragalus Extract 100 mg
Vit A, Vit D3, Vit E, Amino Acids,
Minerals, Probiotic, Polypeptide etc.
Carrier upto 1 g

Indications:

Weight gain fatter promoter, decrease FCR and mortality, growth promoter.
1.Promote nutrition absorption & protein synthesis, promote growth forcefully.
2.Promote bone & visceral organ growth synchronously, shorten breeding cycle. Increase flocks uniformity.
3.Stimulates antibodies production for better immunity, decrease mortality.
4.Decrease internal tract disease, decrease drug input cost.

Dosage&Administration:

For oral administration
Poultry: 1g per 1.5 L water for 4-6 hours drinking.
Use it from 20-25 days before output until output

Withdrawal Period:

None

Storage :

Store in a dry, dark place between 5°C and 25°C.
Keep it away from children.

Packing :

1kg×15 Bags/CTN or 10 Bags/Bucket

Validity :

3 years



Herb Extract





A-HERB ORAL

Herb Extract For Virus Diseases

Herb Extract

For Veterinary Use Only

Composition:

Astragalus polysaccharides $\geq$ 30mg.
Solvents upto 1ml

Indications:

1. Promote immune system growth, Improve animal immunity,.
2. Prevent and treat viral disease such as ND , infectious bursal disease (IBD).
3. Promote to form antibody and improve the activity of macrophage. enhance the antiviral and antibacterial effect.
4. Adjust intestinal function, improve the growth of probiotics, protect intestinal mucosa, increase feed intake and promote growth.

Usage & Dosage:

For Oral Administration

Poultry, Turkey: 1ml per 800-1000ml of drinking water for 3-5 days

Double dosage for serious condition.

Note: Please shake well before use.

Withdrawal Period:

None

Storage :

Keep in cool and shade place, 5-25°C.


Keep out of reach of children.

Packing :

250mL 500ml 1L plastic bottle

Validity :

3 years



GINYE BIOTECH EXPORT

B-HERB ORAL

Immunity Booster & Antivirus Agent Pure Herb Extract Solution

Composition:

Honeysuckle Flower Extract , (Chlorogenic acid $\geq 18\text{mg/mL}$)
Radix Scutellariae Extract (Baicaline $\geq 1.7\text{mg/mL}$)
Solvents upto 1mL

Indications:

It is used for promote immune organs growth, strengthen the immune effect and prevent of vaccination stress.
Prevent and treatment virus / bacterial mixed infection diseases, especially during high-incidence season of virus and bacterial diseases mixed infection, like influenza, newcastle disease, infectious bronchitis, gram bacteria etc.

Dosage&Administration:

For Oral Administration
Poultry, Turkey: 1ml per 750-1000ml of drinking water for 3-5 days
Double dosage for serious condition.
Note: Please shake well before use.

Withdrawal Period:

None

Storage :

Store in dry & cool place, 5-25 °C.
Store in closed packing.

Packing :

250 ml , 1000 ml / plastic bottle

Validity :

3 years

Herb Extract

For Veterinary Use Only





HERB-COC

New Type Herbal Extraction for Coccidiosis

Herb Extract

For Veterinary Use Only

Composition: per ml contains

Purslane Extract.....	20 mg
Poplar Flower Extract.....	20 mg
Areca-nut Extract.....	10 mg
Vitamin K3	10 mg
Solvents upto.....	1 ml

Indications:

- 1.Repair Intestinal Mucosa and restore intestinal function rapidly.
- 2.Used for prevent and treatment coccidiosis of all stages like schizogony and gametogony stages stages of Eimeria spp. in chicken and turkey.
- 3.Make up the defect of traditional coccidium medicine drug resistance problem (such as Toltrazuril , Diclazuril,Sulfonamides), this product no drug resistance for long period medication.
- 4.Pure herbal extract product can be used for all stage of poultry. Green and healthy, no drug residue.

Usage & Dosage:

For oral administration via drinking water:

Poultry : 1ml per 1L-1.5L water, for 3-5 consecutive days.

Double dosage during serious condition.

Notice: This product has better treatment result when use with enteritis medicines.

Withdrawal Period:

None

Storage :

Keep in cool and shade place, 5-25°C.


Keep out of reach of children.

Packing :

Bottle of 250ml 500 ml 1000ml.

Validity :

3 years



GINYE BIOTECH EXPORT

GINHERB WS

Respiratory System Health Defender Herb Extract
Water Soluble Powder

Herb Extract

For Veterinary Use Only



Composition:

Liquorice Extract ≥ 50 mg
Astragalus Extract ≥ 50 mg
Carrier upto 1 g

Indications:

1. Improve animal immunity, protect respiratory tract mucosa, anti-inflammation, anti-infection.
2. Prevent and treatment respiratory tract disease caused by virus and bacterial mixed infection. (such as IB salmonella E.Coli Staphylococcus aureus typhoid bacillus etc)
3. Green healthy, antibiotic free, comply with food safety requirement.

Dosage&Administration:

For Oral Administration

Poultry: 1 g per 1L-1.5L water for poultry drinking 5-8 hours.

Withdrawal Period:

None

Storage :

Store in dry & cool place, 5-25 °C.
Store in closed packing.

Packing :

100g 250g 1kg Bags

Validity :

3 years





GIN-AID WS

Respiratory System Health Defender
Bronchial Embolism Saver

Herb Extract

For Veterinary Use Only

Composition:

Liquorice Extract.....50 mg
Carrier Eucalyptus Extract Etc.

Indications:

- 1.GIN-AID is first-aid product for treatment poultry bronchial embolism, control death rate very soon.
- 2.Protect respiratory tract mucosa, anti-inflammation , anti-infection.
- 3.Prevent and treatment snore bronchial embolism cough caused by virus and bacterial mixed infection. (such as IB salmonella E.Coli Staphylococcus aureus typhoid bacillus etc)
4. Green healthy, antibiotic free, no drug residue.

Dosage&Administration:

For Oral Administration

Poultry: 1 g per 1 liter of water for poultry drinking more than 12 hours.

Double dosage under serious condition.

Withdrawal Period:

None

Storage :

Store in dry & cool place, below 30 °C.
Keep out of reach of children.

Packing :

500g 1kg Bags

Validity :

3 years



GINYE BIOTECH EXPORT

GIN-TOX PLUS

Mold & Bacterial Inhibitor Mycotoxin Cleaner & Immune Booster

Herb Extract

For Veterinary Use Only



Composition:

Each ml contains

Cinnamon Oil 15 mg

Oregano Oil 25 mg

Glycyrrhizic acid.....40 mg

Malic Acid 40 mg

Citric Acid.....55 mg

Glucose Oxidase, Copper Sulfate, Solvents

Indications:

1. Suppression of mycotoxicosis, reduce mortality, improve FCR.
2. Eliminate mycotoxins present in feed and digestive tract, Improve feces consistency and reduce enteric problems.
3. Help repair digestive tract mucosa rapidly, stimulates digestive enzymes production.
4. Reduce farm smell and keep safer environmental condition keep water pipeline cleaning.
5. Long action period, not corrosive, not reduce absorption of nutrients, not alter the taste.

Dosage&Administration:

For oral drinking water:

1L per 1500-2000L Drinking water, keep drink more than 8 hours.

Keep 5-7 days.

Double dosage during mycotoxin serious period.

Storage :

Store in a dry, dark place between 5°C and 25°C.

Keep it away from children.

Packing :

1L 5L 25L Plastic Container.

Validity :

3 years





ANTI-ES W/S

Poultry Formula For E coli Salmonella
Antibiotic-Free W/S

Herb Extract

For Veterinary Use Only

Composition:

Each gram contains

Purslane Extract.....100 mg

Astragalus Extract.....50 mg

Bacillus subtilis $\geq 1 \times 10^{10}$ CFU

Carrier : Potassium Sorbate Starch Yeast Cell wall, etc.

Indications:

This product is new antibiotic-free formula for treatment E-coil Salmonella disease on poultry, its used for preventing and treatment gastrointestinal tract inflammation. it can help animal repair intestines and stomach mucosa rapidly, decrease mortality rate, improve poultry immunity and FCR.

Usage & Dosage:

Normal condition: 1 g per 1-1.5 liter of drinking water 6-8 hours keep 3-5 days.

Serious condition: 1 g per 0.5-1 liter 6-8 hours of drinking water keep 3-5 days.

Withdrawal Period:

None

Storage :

Stored with sealed packaging in a cool, dry and ventilated place.


Avoid to be exposed to the sun,rain,high temperature and high humidity.

Packing :

250g 500g 1kg bags bulk packing forms.

Validity :

3 years



Herb-Care

Liver Kidney Function Booster

Herb Extract

For Veterinary Use Only



Composition:

Eucommia ulmoides Oliv extract, Licorice extract, Eupatorium extract, etc.

Description:

Natural herb extract product to protect poultry's kidney and liver, no residual, no harm, no resistance. Optimize renal function, strengthen liver function.

Indications:

1. Prevent and control all kinds of kidney and liver diseases effectively, such as swollen- kidney, bleeding-kidney, spotted-kidney, hepatitis, fatty liver and etc.
2. Reduce the uric acid salt concentration in the blood, reduce the generation of urate crystals.
3. Treatment and protect damaged liver and kidney caused by drug poisoning or feed poisoning.
4. Improve digestion and metabolism of main feed nutrients especially fats.
5. Support the natural detoxification function of the liver.

Usage & Dosage:

For treatment: 1000ml/500L of water, continuous use for 5-7 days.

For prevention: 1000ml/800L of water, continuous use for 7-10 days.

Storage :

Store in a dry, dark place between 5°C and 25°C.
Keep it away from children.

Packing :

500ml 1000ml per bottle.

Validity :

3 years





RESPIGIN PLUS

Herb Extract

For Veterinary Use Only

Respiratory System Health Defender
Bronchial Embolism Saver

Composition:

Menthol	40 mg
Bromhexine Hcl.....	20 mg
Eucalyptus Oil	10 mg
Solvents upto.....	1 ml

Indications:

- 1.RESPIGIN Plus is first-aid product for treatment poultry bronchial embolism, control death rate very soon.
- 2.Protect respiratory tract mucosa, anti-inflammation, anti-infection, improve immunity rapidly.
- 3.Prevent and treatment snore bronchial embolism cough caused by virus and bacterial mixed infection. (such as IB salmonella E.Coli Staphylococcus aureus typhoid bacillus etc)

Dosage&Administration:

For Oral Administration

Poultry: 1 ml per 1.5-2.0 liter of water for poultry drinking more than 8 hours, keep using 3-5 days. Double dosage under serious condition.

Withdrawal Period:

None

Storage :


Store in dry & cool place, below 30 °C.

Packing :

500ml 1000ml per bottle.

Validity :

3 years



GINYE BIOTECH EXPORT

GINKILL

Fly Mosquito Parasite Killing Agent

Herb Extract

For Veterinary Use Only



Composition:

Each gram contains

Camphor50mg

Cypermethrin50mg

Indications:

Ginkill is new formula long-action & low toxicity farm fly mosquito parasite killer, animal farm environment modifying agent. This product acting very fast, 3-5 mins can see death fly. This product have good result to kill fly mosquito eggs. One time use can be effective 30-60 days.

This product also has good result on killing animal skin parasites such as louse flea and ticks.

Use place :

Animal farm, home, hotel, school, hospital, etc.

Dosage&Administration:

1.Spray: 50g per 10-15L water spray to floor or other surface to moist.

2.Dry Splash: 50g per 200M2 space.

3.Medicated Bath: 50g per10L water wash animal skin or bath wash to fur totally drenched, Reuse it after 7 days.

Note:

1.this product cannot mix use with alkaline substance, use out once prepared.

2.Do not spray to food and feeds or tableware feeding tools

3.This product has toxicity to fish silkworm, donot put packing bag in pond and near water source.

Storage :

Store in a dry, shade place, Keep it away from children.Caked product not affect result.

Packing :

50g/sachet

Validity :

3 years





GINKILL PLUS

Compound Cyromazine WSP

Herb Extract

For Veterinary Use Only

Composition:

Each gram contains

Cyromazine50 mg

Protease, amylase carrier upto 1 g

Indications:

GINKILL PLUS can be used for chicken, laying hens, broilers, the growth of pigs, pregnancy or lactation sow, and other livestock, to control the number of livestock and poultry field fly and larvae, it is conducive to the improvement of livestock and poultry breeding environment, promote the healthy growth of animals and poultry, this product is easy to use, through the drinking water or feed given to control the growth of flies, and do not affect the rate of laying hens, broilers growth and poultry breeding, without a withdrawal period, after the use of livestock and poultry dung will not affect the normal growth of crops, no pollution to the environment.

Character:

- 1.Its water soluble powder, easy to use save time and work.
- 2.Little ammonia in animal house, dry excrement good commercial use value.
- 3.Its not harmful to other animals, no pollution, safe and long acting.
- 4.Little drug residue, safe for food, animal will get little disease brings by fly.

Dosage&Administration:

- 1.Poultry: 100g per 2000L water or 1000kg feed 4-6 weeks continuously, Interval of 4 - 6 weeks, then use 4-6 weeks again with same dosage.
- 2.Livestock: 100g per 1500L water or 750kg feed 4-6 weeks Continuously, Interval of 4 - 6 weeks, then use 4-6 weeks again with same dosage.
- 3.Spray: 100g per 15L water spray to excrement 20m² area.

Storage :

Store in a dry, shade place, Keep it away from children.Caked product not affect result.

Packing :

100g×100sachet/ctn

Validity :

3 years



GINYE BIOTECH EXPORT

SprayCare

Special Treatment Livestock Aphtha / Hoof rotten / Dermatitis / Eczema

Herb Extract

For Veterinary Use Only



Composition:

Butyric acid	30mg
Propionic acid.....	22mg
Plant Soot.....	20mg
TP(tea polyphenol).....	5mg

Indications:

This product is antibiotic free special research for livestock Aphtha / Hoof rotten caused by FMD, Dermatitis / Eczema of piglet, rash trauma, this product is very effective on virus bacterial fungus outside infection treatment.

Usage & Dosage:

1. Aphtha / Hoof rotten, Dermatitis / Eczema: spray 2-3 time per day, and keep 4-5 days.
2. Piglets staphylococcus Dermatitis / Eczema, Circular pustules, 2 time per day, keep 3-4 days.
3. Ruminant Rash, Swine dead skin: 1:5 dilute with water, 2 time per day, keep 5-6 days.
4. Mosquito and fly bites, Reduce swelling and relieve itching: 2 time per day keep 3-4 days.
5. Skin injury, surgical trauma, breast inflammation, 2-3 time per day keep 3-5 days.

Withdraw Period:

None

Storage :

Store in a dry, dark place between 5°C and 25°C.
Keep it away from children.

Packing :

200g 300g 500g bottle.

Validity :

3 years





IMU-C W/S

Poultry Functional Feed Supplement
Super Immune Booster

Nutrition-Powder

For Veterinary Use Only

Composition: Each gram contains

Zymosan	100 mg
Vitamin C	50 mg
Immunoglobulin Yolk (IgY) and Carrier	

Indications:

IMU-C W/S. mainly used for improving animal immunity and prevent or auxiliary treatment of atypical Newcastle disease, mild influenza, adenovirus, immunosuppressive diseases and other viral diseases; precise release of immunosuppressive factors, improve body disease resistance , and quickly restore the sub-health of poultry.

When use it.

Prevention use:

1. Used in the immunization blank period of 16-23 days and 28-35 days of broiler, it can significantly increase the antibody level of Newcastle disease, influenza, and adenovirus, increase resistance, help chickens safely pass through the dangerous period, and avoid and reduce diseases The hazards brought about by the high incidence period.
2. Drinking layer are 30-50 days old, 80-100 days old, and during the peak egg production period can reduce stress, reduce the risk of high incidence of Newcastle disease, influenza, adenovirus and other diseases, and increase breeding benefits.
3. Meat ducks at 10-15 days old and 25-30 days old use this product to increase the antibody levels of paramyxovirus, influenza virus and adenovirus, increase resistance, and avoid and reduce the harm caused by the high incidence of diseases.

Treatment use:

1. It can effectively treat Newcastle disease, influenza, and adenovirus infection in chicken at various stages, quickly and effectively neutralize the diseased chicken virus, increase antibody levels, relieve symptoms, control death, rapidly increase feed intake, and restore the mental state of chicken.
2. It has a significant effect on the paramyxovirus, influenza virus and adenovirus that are popular in the meat duck market. It can quickly control death, increase feed intake, and restore the mental state of large groups.
3. For the treatment of pericardial effusion caused by adenovirus in poultry and waterfowl.

Withdrawal Period:

None

Storage :

Store in a dry, dark place below 5-25°C.
Keep it away from children.

Packing :

500g 1kg bag or jar

Validity :

3 years





Antibiotic





TYLODOX 10%

Tylosin Tartrate 10% + Doxycycline Hcl 10%
Water Soluble Powder

Antibiotic

For Veterinary Use Only

Composition:

Each gram contains

Tylosin Tartrate100 mg

Doxycycline Hydrochloride..... 100 mg

Indications:

Gastrointestinal and respiratory infections caused by tylosin and doxycycline sensitive micro-organisms, like Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus, Streptococcus and Treponema spp. in poultry, calves, goats, sheep and swine.

Dosage&Administration:

For oral administration:

Poultry and swine: 1g per 1 - 2 litres of drinking water for 3 - 5 days

Calves, goats and sheep: Twice daily 1g per 20 kg body weight for 3 - 5 days.

Withdrawal Period:

For meat:

Calves, goats, and sheep : 14 days.

Swine, poultry : 7days.

Storage :


Store in dry dark place between 5°C and 25 °C,
Keep out of reach of children.

Packing :

100g 250g 500g 1kg sachet or plastic jar

Validity :

3 years



GINYE BIOTECH EXPORT

AMOXIGIN 50%

Amoxicilline 50% W/S

Antibiotic

For Veterinary Use Only



Composition:

Each gram contains

Amoxicilline (as Trihydrate).....500 mg

Indications:

AMOXIGIN 50% wsp is effective against infections caused by micro-organisms sensitive to amoxicillin like gastrointestinal infections, respiratory infections, urogenital infections, local inflammations and secondary bacterial infections during viral diseases with cattle, calves, pigs and poultry.

Dosage&Administration:

For oral administration

Poultry : 1g per 4 litres of drinking water during 3 - 5 days.

Cattle, calves : 1 g per 20 kg bodyweight daily, during 3 - 5 days.

Pigs : 1 g per 10 litres of drinking water during 3 - 5 days.

The above mentioned dosages can be doubled without any danger for the animals.

Mixed with feed, the product should be used immediately.

Other dosage should comply veterinarian suggestion.

Medicated drinking water should be used within 24 hours.

Withdrawal Period:

For Meat: 7 days

For Milk : 3 days

For Eggs: 3 days

Storage :

Store in a dry, dark place between 5 °C and 25 °C.

Keep medicine away from children.

Packing :

In a 100g 250g 500g 1kg sachet

Validity :

3 years





AMPROGIN 20%

Amprolium hydrochloride 20% W/S

Antibiotic

For Veterinary Use Only

Composition:

Each gram contains

Amprolium hydrochloride.....200 mg

Indications:

It is used as a preventive or therapeutic agent against Eimeria infections in poultry, especially E. tenella, E. necatrix, E. acervulina and E. praecox.

It is effective against other protozoal infections like Histomoniasis (Blackhead) in turkeys and poultry; against coccidiosis in calves, sheep and goats. against amebiasis in various species.

Dosage&Administration:

For oral administration:

Calves, sheep and goats:

Preventive : 1 g per 40 kg body weight through drinking water or milk for 21 days.

Curative: 1 g per 20 kg body weight through drinking water or milk for 5 days.

Poultry:

Preventive: 1 g per 3 litres of drinking water for 1-2 weeks.

Curative: 1 g per 1.5-2 litres of drinking water for 5-7 days.

Do not use in layers producing eggs for human consumption.

Withdrawal Period:

For meat: 3 days

Storage :


Store in dry dark place between 5°C and 25 °C,
Keep out of reach of children.

Packing :

100g 500g 1kg sachet

Validity :

3 years



GINYE BIOTECH EXPORT

ENROGIN 10%

10% Enrofloxacin Solution

Antibiotic

For Veterinary Use Only



Composition:

Each ml contains 100mg of Enrofloxacin

Indications:

Enrofloxacin oral Solution is a broad spectrum antibacterial for single or mixed bacterial infections, especially for infections caused by anaerobic bacteria. In livestock and canines, Enrofloxacin is effective against a wide range of Gram positive and Gram negative organisms causing infections such as Bronchopneumonia and other respiratory tract infection, gastro enteritis, calf scours, Mastitis, Metritis, Pyometra, Skin and soft tissue infections, ear infections, secondary bacterial infections such as those caused by E.Coli, Salmonella Spp. Pseudomonas, Streptococcus, Bronchiseptica, Klebsiella etc.

Dosage&Administration:

For oral administration:

Poultry: 1 L/1500-2000 L drinking water for 3-5 days.

Swine: 1 L/ 1000-3000 L drinking water for 3-5 days.

Calves, goats and sheep : Twice daily, 10 ml/75 - 150 kg. body weight for 3 - 5 days.

Precaution:

- 1.The product should not be administered at horses and dogs younger than 12 months.
- 2.Avoid direct contact with the product .It is possible to cause dermatitis by contact.
- 3.Overdose can cause digestive disorders as vomit, anorexia, diarrhea and even toxicosis. In that case the administration must be stopped at once and the symptoms must be handled.

Withdrawal Period:

For meat: 8 days

Storage :

Tightly sealed, Store in 5-30°C and avoid light.

Packing :

1L×12Bottles/CTN

Validity :

3 years

GINYE BIOTECH EXPORT





TIMIGIN 25%

25% TILMICOSIN ORAL SOLUTION

Antibiotic

For Veterinary Use Only

Composition:

Contains per ml:

Tilmicosin (as phosphate)..... 250 mg

Indications:

TIMIGIN 25% is indicated for the control and treatment of respiratory infections associated with micro-organisms susceptible to tilmicosin, such as: *Mycoplasma* spp., *Pasteurella multocida*, *Mannheimia haemolytica*, *Actinomyces pyogenes* and *Actinobacillus pleuropneumoniae* in calves and pigs. TIMI ORAL 25% Oral can be used for the treatment of infections in poultry and turkeys caused by *Mycoplasma gallisepticum* (CRD), *Mycoplasma synoviae* (arthritis/synovitis), *Mycoplasma meleagridis* (in turkeys) and by *Clostridium perfringens* (necrotic enteritis).

Dosage&Administration:

For oral administration:

Poultry turkey : 1 ml per 3-4 litres of drinking water (75 ppm) for 3 days.

Calves : Twice daily, 1 ml per 20 kg body weight via (artificial) milk for 3 - 5 days.

Swine : 1 ml per 1-1.5 litres of drinking water (200 ppm) for 5 days.

Withdrawal Period:

Meat: Calves, pigs : 14 days

Poultry, turkeys : 12 day

Storage :

Store in dry dark place between 5°C and 25 °C,

Store in closed packing.

Packing :

In 100ml 250ml and 500ml 1000ml plastic bottle.

Validity :

3 years



GINYE BIOTECH EXPORT



GINCOX 2.5% ORAL

2.5% TOLTRAZURIL ORAL SOLUTION

Antibiotic

For Veterinary Use Only

Composition:

Toltrazuril.....25 mg
Solvents upto.....1 ml

Features:

1. GINCOX 2.5% is used for prevent and treatment of cecum & small intestine coccidiosis.
2. This product can reach coccidium infected position directly, take effect promptly.
3. Newest workmanship and formula and long action, act on all reproduction and grow stage of coccidium.
4. Not easy to get drug resistance.

Indications:

Specially for treatment of Coccidiosis of all stages like schizogony and gametogony stages of Eimeria spp. in chickens and turkeys

- Eimeria acervulina, brunetti, maxima, mitis, necatrix and tenella in chicken.
- Eimeria adenoides, galloparronis and meleagridis in turkey.

Dosage&Administration:

1ml per 1-1.5 Liter of water for poultry drinking freely ,
for 3-5 consecutive days.

Withdrawal Period:

For meat: 5 days

Storage :

Store in dry dark place between 5°C and 25 °C,
Store in closed packing.

Packing :

In 100ml 250ml and 500ml 1000ml plastic bottle.

Validity :

3 years



GINYE BIOTECH EXPORT

FLORGIN 10% ORAL

10% Florfenicol Oral Solution

Antibiotic

For Veterinary Use Only



Composition:

Contains per ml:

Florfenicol..... 100 mg

Excipients ad 1 ml

Indications:

FLORGIN 10% Oral is indicated for preventive and therapeutic treatment of gastrointestinal and respiratory tract infections, caused by florfenicol sensitive micro-organisms such as Actinobaccillus spp. Pasteurella spp. Salmonella spp. and Streptococcus spp. in swine and poultry. The presence of the disease in the herd should be established before preventive treatment. Medication should be initiated promptly when respiratory disease is diagnosed.

Dosage&Administration:

For oral administration. The appropriate final dosage should be based on the daily water consumption.

Poultry : 1 litre per 1000 litres of drinking water (100 ppm; 10 mg/kg body weight) for 3 days.

Swine : 1 litre per 1000 litres of drinking water (100 ppm; 10 mg/kg body weight) for 5 days.

Withdrawal Period:

- For meat:

Poultry : 7 days.

Swine : 21 days.

Storage :

Store in dry dark place between 5°C and 25 °C,
Store in closed packing.

Packing :

100ml and 1000 ml Bottle

Validity :

3 years





ALBEN-100 ORAL

ALBENDAZOLE 10% ORAL SUSPENSION

Antibiotic

For Veterinary Use Only

Composition:

Contains per ml:

Albendazole.....100 mg

Indications:

ALBEN-100 ORAL is indicated for prophylaxis and treatment of worm infections in cattle, calves, sheep and goats like:

Gastrointestinal worms : Bunostomum, Cooperia, Chabertia, Haemonchus, Nematodirus, Oesophagostomum, Ostertagia, Strongyloides and Trichostrongylus spp.

Liver fluke infestations : Adult Fasciola hepatica.

Lungworms : Dictyocaulus viviparus and D. filaria.

Tapeworms : Moniezia spp.

Dosage&Administration:

For oral administration:

Goats and sheep : 1 ml. per 20 kg. body weight.

Calves and cattle : 1 ml. per 12 kg. body weight.

Liver-fluke : 1 ml. per 10 kg. body weight.

In case of heavy infestation or re-infection repeat after 3 - 4 weeks.

Medicated water should be used within 24 hours.

Shake well before use.

Withdrawal Period:

Meat : 12 days

Milk : 4 days

Storage :

Store in dry dark place between 5°C and 25 °C, Store in closed packing.

Packing :

In 500ml 1000ml plastic bottle.

Validity :

3 years



GINYE BIOTECH EXPORT

BROMGIN 1%

Compound Bromhexine Hcl Oral Solution

Antibiotic

For Veterinary Use Only



Composition:

Each ML Contains:

Bromhexine Hcl.....10 mg

Menthol.....20 mg

Indications:

It is a very effective Cough Expectorant. It is indicated To treat symptoms that result From respiratory infections Such as Difficulty in breathing Sneezing in poultry. It will support to prevent influenza type infections.

Dosage&Administration:

1 ml per 3-4 liters of drinking water for 3-5 days.

Withdrawal Period:

Do not use animal products intended

For human consumption during treatment and within 8 days from the last treatment.

Storage :

Keep it in a cool and dry place, protect from the sun light and keep in its original package until the time of its use.

Packing :

500ml 1L

Validity :

3 years





ALBEN-600 BOLUS

Albendazole Bolus 600mg

Antibiotic

For Veterinary Use Only

Composition:

Contains per bolus.:

Albendazole600 mg

Carrier upto.....4.5 g

Description:

Albendazole is a synthetic anthelmintic which belongs to the group of benzimidazole-derivatives with activity against a broad range of worms and at a higher dosage level also against adult stages of liver fluke.

Indications:

Prophylaxis and treatment of worm infections in calves and cattle like:
Gastrointestinal worms : Bunostomum, Cooperia, Chabertia, Haemonchus, Nematodirus, Oesophagostomum, Ostertagia, Strongyloides and Trichostrongylus spp.
Lung worms : Dictyocaulus viviparus and D. filaria.
Tapeworms : Moniezia spp.
Liver fluke : adult Fasciola hepatica.

Contra-indications:

Administration in the first 45 days of gestation.

Side effects: Hypersensitivity reactions.

Dosage&Administration:

For oral administration.

Calves and cattle : 1 bolus per 100 kg body weight.

Sheep and goats : 1 bolus per 50 kg body weight.

For liver fluke : 1 bolus per 50 kg body weight.

Withdrawal Period:

- For meat : 12 days.

- For milk : 4 days.

Storage :

Store in dry dark place between 5°C and 25 °C,
Keep out of reach of children.

Packing :

50 boluses in a box with blister

Validity :

3 years



GINYE BIOTECH EXPORT

ALBEN-2500 BOLUS

Albendazole Bolus 2500mg

Antibiotic

For Veterinary Use Only



Composition:

Contains per bolus.:

Albendazole2500 mg

Carrier upto.....18 g

Description:

Albendazole is a synthetic anthelmintic which belongs to the group of benzimidazole-derivatives with activity against a broad range of worms and at a higher dosage level also against adult stages of liver fluke.

Indications:

Prophylaxis and treatment of worm infections in calves and cattle like:

Gastrointestinal worms : Bunostomum, Cooperia, Chabertia, Haemonchus, Nematodirus, Oesophagostomum, Ostertagia, Strongyloides and Trichostrongylus spp.

Lung worms : Dictyocaulus viviparus and D. filaria.

Tapeworms : Moniezia spp.

Liver fluke : adult Fasciola hepatica.

Contra-indications:

Administration in the first 45 days of gestation.

Side effects: Hypersensitivity reactions.

Dosage&Administration:

For oral administration.

Calves and cattle : 1 bolus per 300 kg body weight.

For liver fluke : 1 bolus per 250 kg body weight.

Withdrawal Period:

- For meat : 12 days.

- For milk : 4 days.

Storage :

Store in dry dark place between 5°C and 25 °C,
Keep out of reach of children.

Packing :

50 boluses in a box with blister

Validity :

3 years





OXYGIN 20%

OXYTETRACYCLINE INJECTION 20% L A W/V

Antibiotic

For Veterinary Use Only

Composition:

Each ml contains oxytetracycline base 200 mg

Indications:

Treatment of diseases caused by oxytetracycline-susceptible organisms in cattle, sheep and goats.

Please see the enclosed insert leaflet for more details.

Dosage&Administration:

For intramuscular or subcutaneous administration:

General: 1 ml per 10 kg body weight.

This dosage can be repeated after 48 hours when necessary.

Withdrawal Period:

28 days for meat and edible tissues

7 days for milk.

Storage :

Store in a dry, dark place between 5 °C and 25 °C. away from direct sunlight.

Keep medicine away from children.

Packing :

In 50ml 100ml glass bottle.

Validity :

3 years



GINYE BIOTECH EXPORT

OXYGIN 5%

OXYTETRACYCLINE INJECTION 5% W/V

Antibiotic

For Veterinary Use Only



Composition:

Each ml contains oxytetracycline.....50 mg

Indications:

Treatment of diseases caused by oxytetracycline-susceptible organisms in cattle, sheep and goats.

Dosage&Administration:

minister by intramuscular injection. The dose is 10mg/kg body weight (2ml/10kg body weight). Treat for 3-5 days depending on the type and severity of the disease.

Cattle	2ml/10kg
Calves	2ml/10kg
Sheep & Goat	2ml/10kg
Pigs	2ml/10kg
Young Animals	4ml/10kg

Withdrawal Period:

21 days for meat and edible tissues 7 days for milk.

Storage :

Store in a cool, dry place under 25 °C, away from direct sunlight.

Packing :

In a 100ml glass bottle.

Validity :

3 years





IVERGIN 1%

IVERMECTIN INJECTION 1% W/V

Antibiotic

For Veterinary Use Only

Composition:

Each ml contains ivermectin10 mg

Indications:

IVERGIN 1% is very effective in the treatment and control of internal and external parasites in cattle, sheep, goats, pigs and camels.

For the treatment and control of gastro-intestinal roundworms, lungworms, grubs, screwworms, fly larvae, lice, ticks and mites in cattle, sheep, goats and camels.

Please see the enclosed insert leaflet for more details.

Dosage&Administration:

For subcutaneous injection only.

Cattle, sheep, goats and camels : 1 ml. per 50kg. bodyweight.

Pigs : 1 ml. per 33kg. bodyweight.

Withdrawal Period:

For meat

Calves cattle, sheep, goats : 28 days.

Swine : 21 days

Storage :

Store in a dry, dark place between

8 °C and 25 °C.

Keep medicine away from children.

Packing :

In 50ml 100ml glass bottle.

Validity :

3 years



GINYE BIOTECH EXPORT

IRONGIN 10%

IRON DEXTRAN INJECTION 10%

Antibiotic

For Veterinary Use Only



Composition:

Each ml contains:

Iron (as iron dextran)100 mg

Vitamin B12.....100 µg

Indications:

IRONGIN 10% Injection is indicated for prevention and treatment of iron deficiency anaemia in piglets, calves, foals, pigs, cattle, horses, sheep, goats and dogs. It is especially necessary in newborns and animals in anaemic conditions.

Dosage&Administration:

For intramuscular or subcutaneous administration:

Calves, foals : 4 - 8 ml in the 1st week of life, when necessary 2nd injection 7 days later.

Piglets : 2 ml on 3rd day of life and on 5th day of life.

Pigs : 3 - 5 ml 1st injection, when necessary 2nd injection 7 days later.

Cattle: 5 - 8 ml 1st injection, when necessary 2nd injection 7 days later.

Sheep, goats : 3 - 4 ml 1st injection, when necessary 2nd injection 7 days later.

Other dosage should comply with veterinarian' s suggestion

Withdrawal Period:

None

Storage :

Store in a cool place between 5°C
25 °C, away from direct sunlight.

Packing :

In a 50ml and 100ml glass bottle.

Validity :

3 years





ENROGIN 10%

ENROFLOXACIN INJECTION 10% W/V

Antibiotic

For Veterinary Use Only

Composition:

Each ml contains Enrofloxacin.....100 mg

Indications:

ENROGIN 10% is indicated against Gastrointestinal and respiratory tract infections caused by enrofloxacin sensitive micro-organisms like Campylobacter, E. coli, Haemophilus, Mycoplasma, Pasteurella and Salmonella spp. in calves, cattle, goats, sheep and swine.

Dosage&Administration:

For intramuscular or subcutaneous administration:

Calves, cattle, sheep and goats : 1 ml per 20 - 40 kg body weight for 3 - 5 days

Swine : 1 ml per 20 - 40 kg body weight for 3 - 5 days.

Withdrawal Period:

For meat:

Calves, cattle, sheep and goats : 21 days.

Swine : 14 days.

For milk : 4 days.

Storage :

Store in a cool place between 5°C- 25 °C,
away from direct sunlight.

Packing :

In 50ml 100ml glass bottle.

Validity :

3 years



GINYE BIOTECH EXPORT

BUTAGIN

BUTAFOSFAN+VITAMIN B12 INJECTION

Antibiotic

For Veterinary Use Only



Composition:

Each ml contains

Butafosfan100 mg

Vitamin B12.....50 µg

Indications:

BUTAGIN Injection is indicated for debilitation by acute or chronic metabolism disorders that result from poor nutrition, inadequate management or disease. Developmental and nutritional disorders in young animals resulting from diseases during the period of growth. As an effective tonic in cases of stress, over-exertion, exhaustion and reduced resistance.

Dosage&Administration:

For subcutaneous, intramuscular or intravenous injection.

Cattle: 5 - 25 ml.

Calves and foals: 5 - 12 ml.

Goats and sheep: 2.5 - 5 ml.

Lambs and kids : 1.5 - 2.5 ml.

Swine: 2.5 - 10 ml.

Piglets: 1 - 2.5 ml.

Repeat daily if required..

Other dosage should comply with veterinarian' s suggestion

Withdrawal Period:

None

Storage :

Store in a cool place between 5°C and 25 °C, away from direct sunlight.

Packing :

In a 50ml and 100ml glass bottle.

Validity :

3 years





DEXAGIN 0.2%

0.2% DEXAMETHASONE INJECTION

Antibiotic

For Veterinary Use Only

Composition:

Each ml contains

Dexamethasone (as Sodium Phosphate).....2 mg

Solvents.....1 ml

Indications:

Dexamethasone is a glucocorticosteroid with a strong antiflogistic, anti allergic and gluconeogenetic action.

Acetone anaemia, allergies, arthritis, bursitis, shock and tendovaginitis in cattle, calves, goats, sheep, swine.

Dosage&Administration:

For subcutaneously and intramuscular administration:

cattle : 5 - 15 ml per animal.

Calves, sheep, goats, pigs : 1 - 2.5 ml per animal.

Other dosage should comply with veterinarian' s suggestion

Withdrawal Period:

Meat : 21 days

Milk: 5days

Storage :

Store in dry dark place between 5°C and 25 °C,

Packing :

In 50ml 100ml glass bottle.

Validity :

3 years



CEFGIN -100

10% Ceftiofur Injectable Suspension

Antibiotic

For Veterinary Use Only



Composition:

Each ml contains

Ceftiofur (as hydrochloride)..... 100 mg

Solvents1 ml

Indications:

CEFGIN 5% is administered to cattle and swine for control of bacterial infections of the respiratory tract, with additional action against foot rot and acute metritis in cattle. It has a wide spectrum of activity against both Gram-positive and Gram-negative bacteria. It exerts its antibacterial action by inhibition of cell wall synthesis.

Dosage&Administration:

For subcutaneously and intramuscular administration:

Cattle: 1 ml per 100 kg body weight for 3 - 5 days, subcutaneously.

Swine: 1 ml per 32 kg body weight for 3 days, intramuscularly.

Shake well before use

Other dosage should comply with veterinarian' s suggestion

Withdrawal Period:

Cattle : 3 days

Swine : 4 days

Storage :

Store in dry dark place between
5°C and 25 °C,

Packing :

In a 50ml and 100ml glass bottle.

Validity :

3 years





FLORGIN 30% INJ.

30% Florfenicol Injection.

Antibiotic

For Veterinary Use Only

Composition:

Contains per ml:

Florfenicol300 mg

Excipients ad1 ml

Indications:

FLORGIN 30% indicated for preventive and therapeutic treatment of respiratory tract infections in cattle due to *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*. The presence of the disease in the herd should be established before preventive treatment. It is additionally indicated for treatment of acute outbreaks of respiratory disease in pigs caused by strains of *Actinobacillus pleuropneumoniae* and *Pasteurella multocida* susceptible to florfenicol.

Dosage&Administration:

For subcutaneous or intramuscular injection.

Cattle:

Treatment (IM) : 1 ml per 15 kg body weight, twice at a 48-h interval.

Treatment (SC) : 2 ml per 15 kg body weight, administered once.

Prevention (SC) : 2 ml per 15 kg body weight, administered once.

Swine : 1 ml per 20 kg body weight (IM), twice at a 48-hour interval.

Withdrawal Period:

- For meat:

Cattle : 30 days (IM route). 44 days

(SC route). Swine : 18 days.

Storage :

Store in a cool place between 5°C and 25 °C, away from direct sunlight.

Packing :

In 50ml 100ml glass bottle.

Validity :

3 years



GINYE BIOTECH EXPORT

TYLOGIN 20%

20% TYLOSIN INJECTION

Antibiotic

For Veterinary Use Only



Composition:

Each 1 ml contains:

Tylosin base.....200 mg

Description:

Tylosin is a macrolide antibiotic with a bacteriostatic action against Gram-positive and Gram-negative bacteria like Campylobacter, Mycoplasma, Pasteurella, Staphylococcus, Streptococcus and Treponema spp.

Indications:

Gastrointestinal and respiratory infections caused by tylosin sensitive micro-organisms, like Campylobacter, Mycoplasma, Pasteurella, Staphylococcus, Streptococcus and Treponema spp. In calves, cattle, goats, sheep and swine.

Contra-indications:

Hypersensitivity to tylosin.

Concurrent administration of penicillins, cephalosporins, quinolones and cycloserine.

Side effects:

After intramuscular administration local reactions can occur, which disappear in a few days. Diarrhoea, epigastric pain and skin sensitization can occur.

Usage and dosage:

For i. m. injection only.

Calves and cows: 1.5-2.5 ml /50kg body weight.

Sheep and goats: 1-3 ml /50kg body weight.

Cats and dogs: 0.1-0.5 ml /10kg body Weight.

(The treatment continues for 3-5days)

Withdrawal Period:

For milk: 3 days. For meat: 5 days.

Storage :

Store in dry dark place between 5°C and 25 °C,

Packing :

In a 50ml and 100ml glass bottle.

Validity :

3 years





IVERGIN SUPER Inj.

IVERMECTIN CLORSULON INJECTION W/V

Antibiotic

For Veterinary Use Only

Composition:

Each ml contains

Ivermectin.....10 mg

Clorsulon 100 mg

Solvents upto.....1 ml

Indications:

Ivermectin is very effective in the treatment and control of internal and external parasites in cattle. Clorsulon is a sulphonamide inactivating adult and immature liver flukes.

IVERCOM SUPER is used for treatment of gastrointestinal roundworms (adults and fourth-stage larvae), lungworms (adults and fourth-stage larvae), liver fluke (*Fasciola hepatica* and *F. gigantica*; adult stages), eye worms, warbles (parasitic stages), sucking lice and mange mites (scabies) in beef cattle and non-lactating dairy cattle.

Dosage&Administration:

For subcutaneous injection only.

Cattle, sheep, goats and camels : 1 ml. per 50kg. bodyweight.

Repeat use should inquire veterinarian.

Withdrawal Period:

For meat : 35 days

Storage :

Store in a dry, dark place between 5 °C and 25 °C.

Keep medicine away from children.

Packing :

Vial of 50 and 100 ml.

Validity :

3 years



100ml

For Veterinary
Use Only

GINYE BIOTECH EXPORT

GINVITA-B

COMPLEX VITAMIN B INJECTION

Antibiotic

For Veterinary Use Only



Composition:

Each ml contains

Vitamin B1.....10 mg

Vitamin B6.....5 mg

D-panthenol.....12.5 mg

Biotin.....100 µg

Solvents ad..... 1 ml

Vitamin B2 5 mg

Vitamin B12.....20 µg

Nicotinamide..... 50 mg

Choline chloride.....10 mg

Indications:

GINVITA-B is a well balanced combination of essential B-vitamins for calves, cattle, goats, horses, sheep and swine. Treatment of B-vitamins deficiencies in farm animals. Improvement of feed conversion and digestion.

Please see the enclosed insert leaflet for more details.

Dosage&Administration:

For subcutaneous or intramuscular administration:

Calves, goats and sheep : 5 - 10 ml.

Lambs piglets : 5 - 8 ml.

Cattle : 10 - 15 ml.

Swine : 2 - 10 ml.

Other dosage should comply with veterinarian' s suggestion

Withdrawal Period:

None

Storage :

Store in a cool place between 5°C and 25 °C, away from direct sunlight.

Packing :

In a 50ml and 100ml glass bottle.

Validity :

3 years





PENSTREP Inj.

Injectable Suspension

Antibiotic

For Veterinary Use Only

Composition:

Contains per ml:

Procaine penicillin G 200 000 IU

Dihydrostreptomycin sulphate 200 mg

Excipients ad 1 ml

Indications:

Arthritis, mastitis and gastrointestinal, respiratory and urinary tract infections caused by penicillin and dihydrostreptomycin sensitive micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Klebsiella, Listeria, Pasteurella, Salmonella, Staphylococcus and Streptococcus spp. in calves, cattle, goats, sheep and swine.

Dosage&Administration:

For intramuscular administration:

Cattle : 1 ml per 20 kg body weight for 3 days.

Calves, goats, sheep and swine : 1 ml per 10 kg body weight for 3 days.

Note: Shake well before use and do not administer more than 20 ml in cattle, more than 10 ml in swine and more than 5 ml in calves, sheep and goats per injection site.

Withdrawal Period:

- For kidney : 45 days.
- For meat : 21 days.
- For milk : 3 days.

Storage :

Store in a dry, dark place between 5 °C and 25 °C.

Keep medicine away from children.

Packing :

Vial of 50 and 100 ml.

Validity :

3 years



GINYE BIOTECH EXPORT

GINVITA Inj.

Multivitamin Injection

Antibiotic

For Veterinary Use Only



Composition:

Contains per ml:

Vitamin A, 10 000 IU
Vitamin E, 20 mg
Vitamin B 2 5 mg
Vitamin B 12 10 µg
Nicotinamide 50 mg
Biotin 125 µg
Excipients ad 1 ml

Vitamin D 3 5 000 IU
Vitamin B 1 10 mg
Vitamin B 6 3 mg
D-panthenol 25 mg
Folic acid 150 µg
Amino acids 12 mg

Indications:

GINVITA Inj. is a well balanced combination of essential vitamins and amino acids for cattle, goats, sheep and swine. It is used for:

Prevention or treatment of vitamin or amino acid deficiencies in farm animals.

Prevention or treatment of stress (caused by vaccination, diseases, transport, high humidity, high temperatures or extreme temperature changes).

Improvement of feed conversion.

Dosage&Administration:

For subcutaneous or intramuscular administration:

Cattle : 10 - 15 ml.

Goats and sheep : 8 - 12 ml.

Swine : 5 - 10 ml.

Withdrawal Period:

None.

Storage :

Store in dry dark place between 5°C and 25 °C,

Keep out of reach of children.

Packing :

50ml and 100ml.

Validity :

3 years





NIXIGIN-34 Inj.

Nitroxynil Injectable Solution

Antibiotic

For Veterinary Use Only

Composition:

Contains per ml:

Nitroxynil340 mg

Indications:

NIXIGIN-34 Inj. is effective against:

Liver fluke infestations caused by *Fasciola hepatica* and *Fasciola gigantica*.
Gastrointestinal parasitism caused by *Haemonchus*, *Oesophagostomum* and *Bunostomum* in cattle, sheep and goats. *Oestrus ovis* in sheep and camels.
Hookworms (*Ancylostoma* and *Uncinaria*) in dogs.

Dosage&Administration:

For subcutaneous or oral administration.

Cattle, sheep, goats, camels : 1 ml per 35 kg b.w. by subcutaneous injection.

This dosage may be increased to 1.25 ml per 35 kg bodyweight in case of acute Fascioliasis (immature flukes).

Dogs : 0.30 ml per 10 kg bodyweight by subcutaneous injection.

No more than 20 ml should be injected into a single site. Repeat treatment only on advice of your veterinarian.

Withdrawal Period:

Meat: Cattle : 60 days

Sheep, goats : 49 days

Storage :

Store in a dry, dark place between 5 °C and 25 °C.

Keep medicine away from children.

Packing :

Vial of 50 and 100 ml.

Validity :

3 years



GINYE BIOTECH EXPORT

AMOXIGIN 15% Inj.

Injectable Suspension

Antibiotic

For Veterinary Use Only



Composition:

Contains per ml:

Amoxycillin (as trihydrate).....150 mg

Solvents ad1 ml

Indications:

Gastrointestinal, respiratory and urinary tract infections caused by amoxycillin sensitive micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, penicillinase negative Staphylococcus and Streptococcus spp. in calves, cattle, goats, sheep and swine.

Contra-indications:

Hypersensitivity to amoxycillin.

Administration to animals with a seriously impaired renal function.

Concurrent administration of tetracyclines, chloramphenicol, macrolides and lincosamides.

Dosage&Administration:

For intramuscular administration:

General: 1 ml per 10 kg body weight, repeatable if necessary after 48 hours.

Shake well before use and do not administer more than 20 ml in cattle, more than 10 ml in swine and more than 5 ml in calves, sheep and goats per injection site.

Withdrawal Period:

- For meat : 21 days.
- For milk : 3 days.

Storage :


Store in dry dark place between 5°C and 25 °C,
Keep out of reach of children.

Packing :

50ml and 100ml.

Validity :

3 years





DIMIGIN PLUS

Sterilized Powder for Injection

Antibiotic

For Veterinary Use Only

Composition:

Contains per sachet of 2.36 gram:

Diminazene aceturate.....1.05 g	Vitamin B6 5 mg
Vitamin B12..... 1 mg	Carrier (Antipyrin) upto.....2.36 g

Contains per sachet of 23.6 gram:

Diminazene aceturate.....10.5 g	Vitamin B6 50 mg
Vitamin B12.....10 mg	Carrier (Antipyrin) upto.....23.6 g

Indications:

Diminazene aceturate belongs to the group of aromatic diamidines which is active against babesiosis, piroplasmosis and trypanosomiasis.

It is used for treatment of babesiosis, piroplasmosis and trypanosomiasis in camels, cattle, goats, sheep and swine horses.

Contra-indications:

Hypersensitivity to diminazene or phenazone.

Administration to animals with an impaired renal and / or hepatic function.

Dosage&Administration:

For deep intramuscular administration:

General: 1 ml per 20 kg body weight (2.36 gram per 300 kg body weight).

Dissolve 2.36 gram powder in 15 ml of sterile water before use.

Withdrawal Period:

- For meat : 21 days.
- For milk : 4 days.

Storage :

Store in dry dark place between 5°C and 25 °C,
Keep out of reach of children.

Packing :

Sachet of 2.36 or 23.6 g 10 sachets/box

Validity :

3 years



5% Closan Inj.

5% Closantel Sodium Injection

Antibiotic

For Veterinary Use Only



Composition:

Contains per ml:

Closantel (as Sodium).....50 mg

Excipients ad1 ml

Indications:

Closantel is a synthetic anti parasitic agent which is highly effective against liver fluke, blood sucking nematodes and larval stages of some arthropods in sheep and cattle. Closantel decouples the mitochondrial oxidative phosphorylation, which leads to the inhibition of ATP synthesis. That causes a significant change in the energy metabolism, and as a consequence the death of the parasite.

Side effects

Hyper acute anaphylactic reactions in cattle.

Hypersensitivity reactions.

Overdoses can cause reduced visibility or blindness, anorexia, lack of coordination and general weakness.

Dosage&Administration:

For subcutaneous administration:

Sheep, cattle less than 200 kg : 1 ml per 10 kg body weight

Cattle over 200 kg : 20 ml plus 5 ml per 100 kg of extra body weight

Do not administer more than 20 ml in cattle per injection site.

Withdrawal Period:

For meat: 50 days.

Storage :

Store in dry dark place between

5°C and 25 °C,

Keep out of reach of children.

Packing :

50ml and 100ml.

Validity :

3 years



GINYE PRODUCTS LIST



Injectable

Composition	Specification	USUAL PACKAGE
Oxytetracycline	5% 10% 20% 30%	
Ivermectin	1% 2%	
Closantel Sodium	5% 10%	
Iron Dextran+B12	10%+100µg /ml	
Complex Vit B	Vit B1 B2 B6 B12	
Enrofloxacin	10% 20%	
Butafosfan+VitB12	100mg+50µg	
Gentamycin Sulfate	5% 10%	
Ceftiofur Hcl	5% 10%	
Dexamethasone Sodium	0.2%	● 10ml
Tylosin	20%	
Florfenicol	10% 20% 30%	● 20ml
Ivermectin+Clorsulon	1%+10%	
Multivitamin	VitA D E B	● 30ml
Procaine penicillin G+Dihydr sulphate	20/20 20/25	
Amoxycillin	15% 20%	
Nitroxynil	25% 34%	● 50ml
VitaminE+Selenium	5%+0.5%	
Diminazene Granular+B6 B12		● 100ml
Tilmicosin Inj.	30%	
Sodium Sulfadiazine +TMP	20%+4%, 40%+8%	● 250ml
Buparvaquone	5%	
Ivermectin+Closantel	1%+5%	● 500ml
Estradiol Benzoate	0.2%	
Lidocaine Hcl	2%	
Oxytocin	10 IU/ml	
Sulfadiazine sodium	33.3%	
Flunixin Meglumine	5%	
Levamisole Hcl	10%	
Ivermectin+Closantel	1%+10%	
Meloxicam	2%	
Amoxicillin + Gentamycin	15%+4%	
Oxytetracycline + FlunixinMeglumine	20%+2%	
Diclofenac sodium	5%	
Vitamin C Inj.	20% 50%	

Oral Solution

Composition	Specification	USUAL PACKAGE
Enrofloxacin Oral	10% 20%	
Tilmicosin	10% 25%	● 100ml
Diclazuril	1% 2.5%	● 250ml
Toltrazuril	2.5% 5%	● 500ml
Toltrazuril+Diclazuril	2.5%+0.5%	● 1L
Ciprofloxacin	10%	● 2.5L
Ivermectin drench	0.08% 0.1%	● 5L
Florfenicol	10% 20% 30%	● 20L
Albendazole Suspension	2.5% 10%	● 25L
Levamisole Hcl+Oxyclozanide	3%+6%	
Albendazole+Ivermectin	2.5%+0.08%	

Powder W/S

Composition	Specification	USUAL PACKAGE
Doxycycline Hcl	10% 20% 50%	
Tylosin Tartrate	10% 50% 100%	
Doxycycline +Tylosin	10%+10%	● 15g
Amoxicillin	10% 50%	● 20g
Colistin Sulfate+Gentamycine	10%+5%	● 50g
Colistin Sulfate+Florfenicol	10%+5%	● 100g
Sulfaclozine Sodium	30%	● 150g
Levamisole Hcl	10% 20%	● 500g
Amprolium Hcl	20% 30%	● 1000g
Dimetridazole+Mequindox	20%+5%	● 5000g
Neomycine Sulphate	20% 50%	
Oxytetracycline+vitamins		
Carbasalate Calcium	50%	
Florfenicol	10% 20% 30%	
Cyromazine	10%	

GINYE PRODUCTS LIST



Herb Extract

Composition	Specification	USUAL PACKAGE
A-Herb Oral	Poultry Virus Disease	
B-Herb Oral	Antivirus immunity booster	● 250ml
Herb-C W/S	Antivirus immunity booster	● 500ml
HERB-COC (Liquid & W/S)	Anti-coccidium oral	● 1L
GIN-HERB	Respiratory System	● 25L
GIN-AID	Bronchial Embolism Snore Cough	● 100g
		● 200g
Menthol Bromhexine Eucalyptus Oil	Respiratory System	● 250g
		● 300g
		● 500g
		● 1kg
Skin Care Spray	Aphtha/Hoof rotten/Dermatitis/Eczema/FMD	● 25kg

Tablets

Composition	Specification	USUAL PACKAGE
Albendazole Bolus	300mg 600mg 2500mg	
Levamisole Bolus	300mg 600mg 1500mg	● 40' s per box
Oxytetracycline Tablet	600mg 1000mg	
Multivitamins Tablet	Vit A D E B Mineral	● 50' s per box
Albendazole + Ivermectin	250mg+5mg	
Tetramisole Hcl	300mg 600mg	● 100' s per box
Levamisole+ Oxyclozanide	300mg+600mg	
Niclosamide	600mg 1250mg	● 250' s per box
Oxfendazole Cobalt Selenium	250mg	
Tylosin	600mg 1000mg	

Nutrition Supplement

Composition	Specification	USUAL PACKAGE
Egg Promoter Oral / Premix	Herb Extract Probiotic	
Vitamins+Herb extract WS		
Multivita WS	Multivita mineral amino acids	● 250ml
Vitamin AD3E WSP		
VITA-C WS	50% 100%	● 500ml
Acidifier liquid	Complex Organic Acid	● 1L
Multivita Solution	Vitamins Amino acids	
Vitamin AD3E Solution	A: 10000IU, D3: 5000IU, E: 20mg	● 5L
Vitamin AD3E Solution	A:20000IU, D3:8000IU, E: 20mg	
Vitamin E+ Selenium	10%+0.15%	● 200g
Complex VitB solution	B1 B2 B6 B12 C K3	
Liver Tonic Liquid	Sorbitol Carnitine etc.	● 250g
Toxin Binder (cleaner) W/S	Yeast Extract Probiotic Potassium sorbate	
Mycotoxin Inhibitor Immune Booster	sodium calcium aluminosilicate Montmorillonite Yeast Cell Wall	● 500g
FAT Growth Promoter	Taurin, Herb extract	● 1kg
Immune Booster W/S	Lysozyme Vit E Vit C Selenium	
IMU-C W/S	Immune booster Antivirus agent	● 25kg
Minerals Premix	All kinds of Mineral	

Disinfectant & Anti-parasite

Composition	Specification	USUAL PACKAGE
Glutaraldehyde+Decamethonium Bromide Compounds	10%+10%	● 50g
Povidone Iodine Solution	10%	● 100g
Glutaraldehyde+Benzalkonium Bromide compounds	10%+10%	● 100ml
Amitraz EC	12.5%	● 1L
Cypermethrin+Camphor (fly kill)	5%+5%	● 5L



Specialized in Animal Health
Care - Offer Service Worldwide

www.ginyechina.com